arXiv:1108.0726v1 [math.PR] 3 Aug 2011

A derivative formula for the free energy function

Yu Zhang

Abstract

We consider bond percolation on the $\mathbf{Z}^d$ lattice. Let M_n be the number of open clusters in $B(n) = [-n,n]^d$. It is well known that $E_p M_n/(2n+1)^d$ converges to the free energy function $\kappa(p)$ at the zero field. In this paper, we show that $\sigma_p^2(M_n)/(2n+1)^d$ converges to $-(p^2(1-p)+p(1-p)^2)\kappa'(p)$.

1 Introduction and statement of results.

Consider bond percolation on the $\mathbf{Z}^d$ lattice, in which bonds are independently open with probability p and closed with probability $1-p$. The corresponding probability measure on the configurations of open and closed bonds is denoted by P_p. We also denote by $E_p(X)$ and $\sigma_p^2(X)$ the expectation and the variance of X with respect to P_p. The open cluster of the vertex x, $\mathbf{C}(x)$, consists of all vertices that are connected to x by an open path. Here an open path from u to v is a sequence $(v_0, b_0, v_1, ..., v_i, b_i, v_{i+1}, ..., v_n)$ with distinct vertices v_i $(0 \leq i \leq n)$ and open bonds b_i adjacent v_i and v_{i+1} such that $v_0 = u$ and $v_n = v$. For vertex set A, A_e denotes the bonds with both vertices in A. Also, $|A|$ denotes the cardinality of A, and $|A_e|$ denotes the number of bonds in A_e. We choose $\mathbf{0}$ as the origin. The percolation probability is

$$\theta(p) = P_p(|\mathbf{C}(\mathbf{0})| = \infty),$$

and the critical probability is

$$p_c = \sup\{p : \theta(p) = 0\}.$$

We denote the open cluster distribution by

$$\theta_n(p) = P_p(|\mathbf{C}(\mathbf{0})| = n).$$

By analogy with the Ising model, we introduce the magnetization function as

$$M(p,h) = 1 - \sum_{n=0}^{\infty} \theta_n(p) e^{-nh} \text{ for } h \geq 0.$$

By setting $h = 0$ in the magnetization function,

$$M(p,0) = \theta(p).$$

AMS classification: 60K35.

Key words and phrases: percolation, free energy, variance.

Using term by term differentiation, we also have

$$\lim_{h\to 0^+} \frac{\partial M(p,h)}{\partial h} = E_p(|\mathbf{C}(\mathbf{0})|; |\mathbf{C}(\mathbf{0})| < \infty) = \chi^f(p).$$

$\chi^f(p)$ is called the *mean cluster size.* The *free energy* $F(p,h)$ is defined by

$$F(p,h) = h(1-\theta_0(p)) + \sum_{n=1}^{\infty} \frac{1}{n}\theta_n(p)e^{-hn} \text{ for } h > 0.$$

If we differentiate with respect to h, then we find

$$\frac{\partial F(p,h)}{\partial h} = M(p,h).$$

For $h > 0$, the free energy is infinitely differentiable with respect to p. If $h = 0$, $F(p,0)$ is called the *zero-field* free energy. The zero-field free energy $F(p,0)$ is a more interesting and more difficult object of study since it is believed that there is a singularity point at p_c. By our definition,

$$F(p,0) = E(|\mathbf{C}(\mathbf{0})|^{-1}; |\mathbf{C}(\mathbf{0})| > 0).$$

Grimmett (1981) discovered that the zero-field free energy also coincides with the number of open clusters per vertex. Let us define the number of open clusters per vertex as follows. We denote by M_n the number of open clusters in $B(n)$. By a standard ergodic theorem (see Theorem 4.2 in Grimmett (1989)), the limit

$$\lim_{n\to\infty} \frac{1}{|B(n)|} M_n = \lim_{n\to\infty} \frac{1}{(2n+1)^d} M_n = \kappa(p) \text{ a.s. and } L_1 \tag{1.1}$$

exists for all $0 \le p \le 1$. $\kappa(p)$ is called the *number of open clusters per vertex.* Grimmett (1981) proved that

$$\kappa(p) = F(p,0). \tag{1.2}$$

$\kappa(p)$, as a function of p, is analytic for $p \neq p_c$ and differentiable on $[0,1]$ (see Kesten (1982)). In particular, $\kappa(p)$ is proved (see Kesten (1982)) to be twice differentiable at p_c for the square lattice. In general, physicists believe that the zero-field free energy is twice, but not three times differentiable at p_c.

On the other hand, for $\kappa(p)$, as a limit of random variables, Zhang (2001) showed the following central limit theorem. For $p \in [0,1]$,

$$\frac{M_n - E_p M_n}{\sigma_p(M_n)} \Rightarrow \text{ a standard normal distribution.} \tag{1.3}$$

Zhang also showed large deviations for M_n. In this paper, we show another property for $\kappa(p)$.

Theorem. *For* $0 \le p \le 1$,

$$\lim_{n\to\infty} \sigma_p^2(M_n)/(2n+1)^d = -(p^2(1-p) + p(1-p)^2)\kappa'(p).$$

Remark. If b_0 is the bond with vertices $\mathbf{0}$ and $(1,0,\cdots,0)$, and $\mathcal{G}(b)$ is the event that there does not exist an open path in $\mathbf{Z}_e^d \setminus b$ connecting the two vertices of bond b, then by (2.9) in the proof of the theorem, we have

$$\kappa'(p) = -dP_p(\mathcal{G}(b_0)). \tag{1.5}$$

By using Sykes and Essam's formula (1964), we know that for the square lattice,

$$\kappa'(0.5) = -1.$$

Thus

$$\lim_{n\to\infty} \sigma_{0.5}^2(M_n)/(2n+1)^d = 0.25. \tag{1.6}$$

2 Proof of theorem.

For any bond b, let $v_1(b)$ and $v_2(b)$ be the two vertices of b. Given p, we start by taking the derivative of $E_p(M_n)$. Note that

$$E_pM_n = \sum_{l=1}^{\infty} P_p(M_n \geq l),$$

and the event $\{M_n \geq l\}$ is decreasing. Let $\{M_n \geq l\}(b)$ be the event that b is a pivotal bond for $\{M_n \geq l\}$. By Russo's formula, note that $P_p(M_n \geq 1) = 1$, so

$$\begin{aligned}
\frac{dE_pM_n}{dp} &= -\sum_{l=2}^{\infty} \sum_{b\in B_e(n)} P_p(\{M_n \geq l\}(b)) \\
&= -\sum_{b\in B_e(n)} \sum_{l=2}^{\infty} P_p(M_n = l-1 \text{ or } l \text{ if } b \text{ is open or closed}) \\
&= -\sum_{b\in B_e(n)} P_p\left(\bigcup_{l=2}^{\infty} M_n = l-1 \text{ or } l \text{ if } b \text{ is open or closed}\right). \qquad (2.1)
\end{aligned}$$

Let

$$\mathcal{E}_n(b) = \{b \text{ is a pivotal bond for the open connection of } v_1(b) \text{ and } v_2(b) \text{ in } B(n)\}.$$

In other words, if b is open, then $v_1(b)$ and $v_2(b)$ are connected by open paths. Conversely, if b is closed, then $v_1(b)$ and $v_2(b)$ are not connected by open paths. Thus, for each $b \in B_e(n)$,

$$P_p\left(\bigcup_{l=2}^{\infty} M_n = l-1 \text{ or } l \text{ if } b \text{ is open or closed}\right) = P_p(\mathcal{E}_n(b)). \tag{2.2}$$

Let $\mathcal{G}_n(b)$ be the event that there does not exist an open path connecting $v_1(b)$ to $v_2(b)$ inside $B_e(n) \setminus b$. Then we would have

$$\mathcal{E}_n(b) = \mathcal{G}_n(b). \tag{2.3}$$



Figure 1: The figure shows that $\mathcal{G}_n(b)$ occurs, but $\mathcal{G}(b)$ does not occur. The two vertices of b are connected by open paths, but they have to reach to the boundary of $B(n)$ before connecting with each other. The dotted path indicates the closed bonds that block the connection of open paths from $v_1(b)$ and $v_2(b)$ inside $B(n)$.

To see (2.3), if there were such a path, then $v_1(b)$ and $v_2(b)$ would always be connected by an open path whenever b is open or closed. So b would not be a pivotal bond. On the other hand, if there does not exist an open path connecting $v_1(b)$ and $v_2(b)$ in $B_e(n) \setminus b$, then b should be a pivotal bond for the open connection of $v_1(b)$ and $v_2(b)$. With these observations,

$$\frac{dE_p M_n}{dp} = -\sum_{b \in B_e(n)} P_p(\mathcal{G}_n(b)) = -\sum_{b \in B(n)} P_p(\mathcal{G}(b)) - \sum_{b \in B_e(n)} P_p\left(\mathcal{G}_n(b) \cap \mathcal{G}^C(b)\right).$$

By translation invariance, the ratio of the first term above is

$$\sum_{b \in B_e(n)} P_p(\mathcal{G}(b))/|B_e(n)| = P_p(\mathcal{G}(b_0)). \tag{2.4}$$

We use the following lemma to estimate the second term.

Lemma.

$$\lim_{n\to\infty} \frac{1}{|B_e(n)|} \sum_{b\in B_e(n)} P_p\left(\mathcal{G}_n(b) \cap \mathcal{G}^C(b)\right) = 0 \textit{ uniformly on } [0,1].$$

Proof. Let $A(n,m) = B(n)\setminus B(m)$ for $m \leq n$, and let $\bar{A}(n,m)$ be the closure of $A(n,m)$. Then

$$\begin{aligned}
& \frac{1}{|B_e(n)|} \sum_{b\in B_e(n)} P_p\left(\mathcal{G}_n(b) \cap \mathcal{G}^C(b)\right) \\
\leq\ & \frac{1}{|B_e(n)|} \sum_{b\in B_e(n-\sqrt{n})} P_p\left(\mathcal{G}_n(b) \cap \mathcal{G}^C(b)\right) + \frac{1}{|B_e(n)|} \sum_{b\in \bar{A}_e(n,n-\sqrt{n})} P_p\left(\mathcal{G}_n(b) \cap \mathcal{G}^C(b)\right) \\
\leq\ & \frac{1}{|B_e(n)|} \sum_{b\in B_e(n-\sqrt{n})} P_p\left(\mathcal{G}_n(b) \cap \mathcal{G}^C(b)\right) + O\left(n^{-0.5}\right), \qquad (2.5)
\end{aligned}$$

where we may assume that $n-\sqrt{n}$ is an integer; otherwise we may use $\lceil n-\sqrt{n}\ \rceil$ to replace $n-\sqrt{n}$. Let us estimate the first term in the above inequality. For $b \in B_e(n-\sqrt{n})$, $\mathcal{E}_n(b) \cap \mathcal{E}^C(b)$ implies that there exists an open path from $v_1(b)$ and $v_2(b)$ without using b, but open paths cannot stay inside $B(n)$. In other words, the open path adjacent to $v_1(b)$ has to reach the boundary of $B(n)$ before reaching $v_2(b)$ (see Fig. 1). Similarly, the open path adjacent to $v_2(b)$ has to reach the boundary of $B(n)$ before reaching $v_1(b)$ (see Fig. 1). Therefore, there exist two disjoint open paths from $v_1(b)$ and $v_2(b)$ such that both reach to the boundary of $v_1(b) + B(\sqrt{n}-1)$. Let $\mathcal{D}(b,\sqrt{n})$ be the event. By using the estimate of Theorem 6.1 in Grimmett (1989), for all $p \in [0,1]$, there exist constant $C = C(d)$ and $\delta = \delta(d) \leq 0.5$ such that

$$P_p\left(\mathcal{D}(b,\sqrt{n})\right) \leq Cn^{-\delta}. \qquad (2.6)$$

By (2.5) and (2.6),

$$\frac{1}{|B_e(n)|} \sum_{b\in B_e(n)} P_p\left(\mathcal{G}(b) \cap \mathcal{G}_n^C(b)\right) \leq O(n^{-\delta}). \qquad (2.7)$$

Therefore, the lemma follows from (2.7). □

It follows from (2.3), the lemma, and (2.4) that

$$\frac{dE_pM_n}{|B(n)|dp} = -(|B_e(n)|/|B(n)|) \sum_{b\in B(n)} P_p(\mathcal{G}(b))/|B_e(n)| + O(n^{-\delta}). \qquad (2.8)$$

Note that

$$|B_e(n)| = 2dn(2n+1)^{d-1} \text{ and } |B(n)| = (2n+1)^d,$$

so if we let $n \to \infty$ in (2.8),

$$\kappa'(p) = -\lim_{n\to\infty} \sum_{b\in B_e(n)} P_p(\mathcal{G}_n(b))/(2n+1)^d = -dP_p\left(\mathcal{G}(b_0)\right). \qquad (2.9)$$

Now we estimate the variance of M_n. We list the bonds in $B_e(n)$ in some order:

$$\{b_1, \cdots, b_k\}.$$

We define the independent Bernoulli-random variables $\{\omega(b_i)\}$ for $1 \leq i \leq k$ to be $\omega(b_i) = 0$ (open) or $\omega(b_i) = 1$ (closed) with probability p or $1-p$. Now we construct the following filtration:

$$\mathcal{F}_0 = \{\emptyset, \Omega\} \subset \mathcal{F}_1 = \{\sigma\text{−field generated by } \omega(e_1)\} \subset, \cdots, \subset \mathcal{F}_k = \{\sigma\text{−field generated by } \omega(e_1), \cdots, \omega(e_k)\}.$$

The martingale representation of $M_n - E_p M_n$ is

$$M_n - E_p(M_n) = \sum_{i=0}^{k-1} [E_p(M_n | \mathcal{F}_{i+1}) - E_p(M_n | \mathcal{F}_i)].$$

Let the martingale difference be

$$\Delta_{i,k} = [E_p(M_n | \mathcal{F}_{i+1}) - E_p(M_n | \mathcal{F}_i)].$$

The variance is

$$\sigma_p^2(M_n) = \sum_{i=0}^{k-1} E_p(\Delta_{i,k}^2). \tag{2.10}$$

Both M_n and $\Delta_{i,k}$ can be viewed as functions on $[0,1]^k$ and $[0,1]^i$, respectively. So we can write $M_n(c_1, \cdots, c_k)$ and $\Delta_{i,k}(c_1, \cdots, c_i)$ for them, where c_i only takes a value of zero or one. Thus

$$\begin{aligned}
& \Delta_{i,k} = \Delta_{i,k}(c_1, \cdots, c_i) \\
= & \sum_{c_{i+1}, \cdots, c_k} M_n(c_1, \cdots, c_i, c_{i+1}, \cdots, c_k) P_p\left(\omega(b_{i+1}) = c_{i+1}, \cdots, \omega(b_k) = c_k\right) \\
& - \sum_{c_i', c_{i+1}, \cdots, c_k} M_n(c_1, \cdots, c_{i-1}, c_i', c_{i+1}, \cdots, c_k) P_p(\omega(b_i) = c_i', \omega(b_{i+1}) = c_{i+1}, \cdots, \omega(b_k) = c_k) \\
= & \sum_{c_i', c_{i+1}, \cdots, c_k} [M_n(c_1, \cdots, c_i, \cdots, c_k) - M_n(c_1, \cdots, c_{i-1}, c_i', c_{i+1}, \cdots, c_k)] \\
& \quad \cdot P_p(\omega(b_i) = c_i', \omega(b_{i+1}) = c_{i+1}, \cdots, \omega(b_k) = c_k),
\end{aligned} \tag{2.11}$$

where the sum takes over all possible values of c_i and c_i'. On $\mathcal{G}_n^C(b_i)$,

$$\begin{aligned}
M_n(c_{i+1}, \cdots, c_i, \cdots, c_k) - M_n(c_{i+1}, \cdots, c_i' = 0, \cdots, c_k) = 0 \text{ for } c_i = 0 \text{ or } c_i = 1, \\
M_n(c_{i+1}, \cdots, c_i, \cdots, c_k) - M_n(c_{i+1}, \cdots, c_i' = 1, \cdots, c_k) = 0 \text{ for } c_i = 0 \text{ or } c_i = 1.
\end{aligned}$$

Thus, by (2.11),

$$\Delta_{i,k}\left(1 - I_{\mathcal{G}_n(b_i)}\right) = 0, \tag{2.12}$$

where $I_{\mathcal{A}}$ is the indicator of $\mathcal{A}$. With this observation,

$$\Delta_{i,k} = \Delta_{i,k} I_{\mathcal{G}_n(b_i)}. \tag{2.13}$$

Note that on $\mathcal{G}_n(b_i)$,

$$[M_n(c_1,\cdots,c_i=1,\cdots,c_k)-M_n(c_1,\cdots,c_{i-1},c_i'=0,c_{i+1},\cdots,c_k)]=-1.$$

so

$$\begin{aligned}
&\Delta_{i,k}(c_1,\cdots,c_{i-1},1)I_{\mathcal{G}_n(b_i)}\\
=\;& I_{\mathcal{G}_n(b_i)}\sum_{c_{i+1},\cdots,c_k}[M_n(c_1,\cdots,c_i=1,\cdots,c_k)-M_n(c_1,\cdots,c_{i-1},c_i'=0,c_{i+1},\cdots,c_k)]\\
&\quad\cdot P_p(\omega(b_i)=c_i'=0,\omega(b_{i+1})=c_{i+1},\cdots,\omega(b_k)=c_k)\\
&+I_{\mathcal{G}_n(b_i)}\sum_{c_{i+1},\cdots,c_k}[M_n(c_1,\cdots,c_i=1,\cdots,c_k)-M_n(c_1,\cdots,c_{i-1},c_i'=1,c_{i+1},\cdots,c_k)]\\
&\quad\cdot P_p(\omega(b_i)=c_i'=1,\omega(b_{i+1})=c_{i+1},\cdots,\omega(b_k)=c_k)\\
=\;& I_{\mathcal{G}_n(b_i)}\sum_{c_{i+1},\cdots,c_k}[M_n(c_1,\cdots,c_i=1,\cdots,c_k)-M_n(c_1,\cdots,c_{i-1},c_i'=0,c_{i+1},\cdots,c_k)]\\
&\quad\cdot P_p(\omega(b_i)=c_i'=0)P_p(\omega(b_{i+1})=c_{i+1},\cdots,\omega(b_k)=c_k)\\
=\;& -(1-p)I_{\mathcal{G}_n(b_i)}
\end{aligned}$$

Therefore,

$$\Delta_{i,k}(c_1,\cdots,c_{i-1},1)I_{\mathcal{G}_n(b_i)}=(1-p)^2 I_{\mathcal{G}_n(b_i)}. \tag{2.14}$$

Similarly, note that on $\mathcal{G}_n(b_i)$,

$$[M_n(c_1,\cdots,c_i=0,\cdots,c_k)-M_n(c_1,\cdots,c_{i-1},c_i'=1,c_{i+1},\cdots,c_k)]=1.$$

so

$$\begin{aligned}
&\Delta_{i,k}(c_1,\cdots,c_{i-1},0)I_{\mathcal{G}_n(b_i)}\\
=\;& I_{\mathcal{G}_n(b_i)}\sum_{c_{i+1},\cdots,c_k}[M_n(c_1,\cdots,c_i=0,\cdots,c_k)-M_n(c_1,\cdots,c_{i-1},c_i'=0,c_{i+1},\cdots,c_k)]\\
&\quad\cdot P_p(\omega(b_i)=c_i'=0,\omega(b_{i+1})=c_{i+1},\cdots,\omega(b_k)=c_k)\\
&+I_{\mathcal{G}_n(b_i)}\sum_{c_{i+1},\cdots,c_k}[M_n(c_1,\cdots,c_i=0,\cdots,c_k)-M_n(c_1,\cdots,c_{i-1},c_i'=1,c_{i+1},\cdots,c_k)]\\
&\quad\cdot P_p(\omega(b_i)=c_i'=1,\omega(b_{i+1})=c_{i+1},\cdots,\omega(b_k)=c_k)\\
=\;& I_{\mathcal{G}_n(b_i)}\sum_{c_{i+1},\cdots,c_k}[M_n(c_1,\cdots,c_i=0,\cdots,c_k)-M_n(c_1,\cdots,c_{i-1},c_i'=1,c_{i+1},\cdots,c_k)]\\
&\quad\cdot P_p(\omega(b_i)=c_i'=1)P_p(\omega(b_{i+1})=c_{i+1},\cdots,\omega(b_k)=c_k)\\
=\;& pI_{\mathcal{G}_n(b_i)}
\end{aligned}$$

Therefore,

$$\Delta_{i,k}(c_1,\cdots,c_{i-1},0)I_{\mathcal{G}_n(b_i)}=p^2 I_{\mathcal{G}_n(b_i)}. \tag{2.15}$$

Together with (2.13), (2.14) and (2.15),

$$\begin{aligned}
E_p\Delta_{i,k}^2 &= E_p\Delta_{i,k}^2 I_{\mathcal{G}_n(b_i)} \\
&= E_p\Delta_{i,k}^2(c_1,\cdots,c_{i-1},1)I_{\{\omega(b_i)=1\}}I_{\mathcal{G}_n(b_i)} + E_p\Delta_{i,k}^2(c_1,\cdots,c_{i-1},0)I_{\{\omega(b_i)=0\}}I_{\mathcal{G}_n(b_i)} \\
&= E_p(1-p)^2 I_{\{\omega(b_i)=1\}}I_{\mathcal{G}_n(b_i)} + E_p p^2 I_{\{\omega(b_i)=0\}}I_{\mathcal{G}_n(b_i)}.
\end{aligned}$$

Note that $\{\omega(b_i)\}$ and $I_{\mathcal{G}_n(b_i)}$ are independent, so

$$E_p\Delta_{i,k}^2 = [(1-p)^2p + p^2(1-p)]E_pI_{\mathcal{G}_n(b_i)} = [(1-p)^2p + p^2(1-p)]P_p(\mathcal{G}_n(b_i)). \tag{2.16}$$

By (2.10) and (2.16),

$$\sigma_p^2(M_n) = \sum_{i=1}^{k} E_p(\Delta_{i,k}^2) = [(1-p)^2p + p^2(1-p)] \sum_{b\in B_e(n)} P_p(\mathcal{G}_n(b)). \tag{2.17}$$

Therefore, if we divide the both sides of (2.17) by $(2n+1)^d$ and let n go to infinity, by (2.9),

$$\begin{aligned}
\lim_{n\to\infty} \sigma_p^2(M_n)/(2n+1)^d &= \lim_{n\to\infty}[(1-p)^2p + p^2(1-p)] \sum_{b\in B_e(n)} P_p(\mathcal{G}_n(b))/(2n+1)^d \\
&= -[(1-p)^2p + p^2(1-p)]\kappa'(p).
\end{aligned}$$

So the theorem follows.

References

Grimmett, G. (1981). On the differentiability of the number of clusters per vertex in the percolation model. *J. Lond. Math. Soc.* **23** 372–384.
Grimmett, G. (1989). *Percolation.* Springer-Verlag, New York.
Kesten, H. (1982). *Percolation Theory for Mathematicians.* Birkhauser, Boston.
Sykes, M. F., and Essam, J. W. (1964). Exact critical percolation probabilities for site and bond problems in two dimensions. *J. Math. Phys.* **5** 1117-1127.
Zhang, Y. (2001). A martingale approach in the study of percolation clusters on the Z^d lattice. *J. Theore. Probab.* **14** 165–187.

Yu Zhang
Department of Mathematics
University of Colorado
Colorado Springs, CO 80933
yzhang3@uccs.edu